

# Oral Pharmaceuticals in Primary Care Optometry

Blair Lonsberry, MS, OD, MEd., FAAO  
 Professor of Optometry  
 Pacific University College of Optometry  
 blonsberry@pacificu.edu

## Disclosures:

- Sun Pharmaceuticals: speakers bureau,
- Dompe: advisory board,
- RVL Pharmaceuticals: advisory board


1

2



## Case

- 20 year old male presents with a red painful eye
  - Started that morning when he woke up
  - reports a watery discharge, no itching, and is not a contact lens wearer
- SLE:
  - See attached image with NaFl stain



3

## Herpes Simplex Virus (HSV) Keratitis: Clinical Features

- Characterized by primary outbreak and subsequent reactivation
  - Primary outbreak is typically mild or subclinical (90% of people are asymptomatic)
  - Most clinical ocular infections are manifestations of virus reactivation; ocular involvement occurs in fewer than 5% of primary infections
- After primary infection, the virus becomes latent in the trigeminal ganglion or cornea
  - The majority of ophthalmic HSV cases are unilateral, with recurrences affecting the same eye. Bilateral disease (not necessarily concurrent) occurs in 1-12% of cases and is more common in patients with atopy or other immune abnormalities
- Stress, UV radiation, and hormonal changes can reactivate the virus
- Lesions are common in the immunocompromised (i.e. recent organ transplant or HIV patients)



3



4

## Herpes Simplex Virus Keratitis

- **Epithelial Keratitis:**
  - Symptoms:
    - Ocular irritation, redness, photophobia, watering, blurred vision
  - Signs:
    - Swollen opaque epithelial cells arranged in a course punctate or stellate pattern
    - Central desquamation results in a dendrite\*\*\*
      1. Central ulceration
      2. Terminal end bulbs
    - \*\*\*Corneal sensation is reduced\*\*\*



## Dendritic Ulcers




6

5

6

### HSV Geographic Ulcer



[https://www.researchgate.net/figure/Geographic-corneal-ulcer-caused-by-herpes-simplex-virus-keratitis\\_fig1\\_26720111](https://www.researchgate.net/figure/Geographic-corneal-ulcer-caused-by-herpes-simplex-virus-keratitis_fig1_26720111)



7

### Pediatric HSV Keratitis

- pediatric herpes simplex keratitis has an 80% risk of recurrence, a 75% risk of stromal disease, and a 30% rate of misdiagnosis
- 80% of children with herpes simplex keratitis develop scarring, mostly in the central cornea
  - results in the development of astigmatism
  - 25% of children have more than 2 D of astigmatism, most of which is irregular
- consider pediatric HSV when a patient has unilateral recurrent disease in the anterior segment



8


### Herpes Simplex Virus Keratitis Management

- Topical:
  - Viroptic (trifluridine) q 2h until epi healed then taper down for 10-14 days.
    - Viroptic is toxic to the cornea.
  - Zirgan (ganciclovir) available, use 5 times a day until epi healed then 3 times for a week (US only)



9

### First pass metabolism



10

10

### Anti-Viral Medication

Drug	Mechanism of Action	Bioavailability	Dosing	Side Effects
Acyclovir	Acyclovir interferes with DNA synthesis inhibiting viral replication	10-30% gets absorbed Short ½ life *Metabolized in kidneys	Simplex: 400 mg 5x/day Zoster: 800 mg 5x/day	Overall very safe Nausea, vomiting, headaches, dizziness, confusion
Valacyclovir	Acyclovir pro-drug Equivalent to acyclovir but better for pain management	95% converted to acyclovir* Better bioavailability and longer 1/2 life	Simplex: 500 mg tid Zoster: 1 g tid	Same as acyclovir
Famciclovir	Inhibits DNA chain elongation It is metabolized to penciclovir where it is active 10-20x as long as acyclovir	Superior to acyclovir*	Simplex: 250 mgTID Zoster: 500 mg TID	Same as acyclovir

11

### HSV Stromal Disease



- HSV Stromal disease is an immune-mediated disease
  - Stromal involvement is rarely an initial ocular finding, accounting for fewer than 2% of initial presentations but for 20 – 60% of recurrent corneal disease
- Increased risk of scarring and high risk of poor visual prognosis
- Requires corticosteroids (HEDS: corticosteroid reduced risk of progression by 68%)
  - Without epithelial defect: corticosteroids and prophylactic anti-viral dosage
  - With epithelial defect: active infection anti-viral dosage with judicious corticosteroids



12

### How much to dose steroid?

- HEDS used QID of *prednisolone phosphate*
- Current Recommendations:
  - Mod – severe (especially with neo): 1% Prednisolone or Lotemax QID to 6x/day
  - Want the lowest dose needed to control the inflammation
  - AAO EBM Treatment Guideline 2014
    - Topical steroid for 10 weeks (this is based on HEDS results) with oral antiviral



13

### HSV Epithelial Keratitis

- Treatment Regimen:
  - Zigan (ganciclovir) available, use 5 times a day until epi healed then 3 times for a week OR
  - Oral Valtrex 500 mg 3x/day for 7-10 days
  - Artificial tears
- L-Lysine 2 grams daily?
  - Proven to "slow down" and retard the growth of the herpes virus and inhibit viral replication
- Debride the ulcer?
  - Prior to topical antiviral therapy debridement was treatment of choice
  - Generally try to avoid use of sharp instruments and use of cotton swab and anesthetic
- RTC 1 day, 4 days, 7 days



14

### Herpes Simplex Keratitis

- Prophylactic Treatment:
  - Reduces the rate of recurrence of epithelial and stromal keratitis by ≈ 50%
    - Acyclovir 400 mg BID
    - Valtrex 500 mg QD
    - Famvir 250 mg QD
  - L-lysine 1 gram/day:
    - Proven to "slow down" and retard the growth of the herpes virus and inhibit viral replication
  - Frequent debilitating recurrences, bilateral involvement, or HSV infection in a monocular patient



15

Academic Emergency Medicine  
ORIGINAL CONTRIBUTION

**Topical Tetracaine Used for 24 Hours Is Safe and Rated Highly Effective by Patients for the Treatment of Pain Caused by Corneal Abrasions: A Double-blind, Randomized Clinical Trial**

Ned Washien, MD, FACEM, Ian K. Dennis, and Peter Herkles, DSc

**Abstract**  
Objective: The objective of this study was to test the hypothesis that topical tetracaine would be safe to use for 24 hours and would not affect corneal healing. This approach would potentially spare pain relief and the associated costs of care in an acute care setting. Study Design: Prospective, randomized trial of tetracaine versus saline on the emergency department (ED) of a regional tertiary care teaching hospital. A total of 100 patients presenting with corneal abrasions were randomized to tetracaine or saline. The intervention was tetracaine, given over the 24-hour period. Patients were blinded to the treatment. The primary endpoint was patient-rated pain relief at 24 hours. Main study endpoint measures were patient ED satisfaction at 48 hours and tetracaine safety and side effects. Secondary endpoints were patient satisfaction at 48 hours and tetracaine safety. Results: Corneal abrasions were treated with topical tetracaine (TET) or saline (SAL) over 24 hours, which results for 48 hours and patient-reported corneal effectiveness using a numeric rating scale (NRS) of 0 to 10, returned during telephone interviews. Results: As hypothesized, tetracaine was superior to saline in all patients in the tetracaine group, and the difference probably confers a net benefit. ED of this scenario is 0.51. There was no significant difference in corneal healing or treatment of patients with corneal abrasions. Conclusions: Topical tetracaine used for 24 hours is safe and while there was no significant difference in patient ED satisfaction at 48 hours, patient ratings of corneal effectiveness showed that patients perceived tetracaine to be significantly more effective than saline. ACADEMIC EMERGENCY MEDICINE 2012,15(1):10-16 © 2012 by the Society for Academic Emergency Medicine

16

### Systemic NSAID's

- NSAID's are the drug of choice for treating mild to moderate ocular pain
  - Very beneficial for treating systemic inflammation as well.
- All NSAID's are rapidly absorbed from the GI tract, highly bound in the plasma, and capable of crossing the blood-brain barrier.
- Exhibit a "ceiling effect" – there is a dosage beyond which no further analgesia occurs
  - Produce no tolerance or dependence, increasing their safety profile.
- Variability exists in patient responses to NSAID's
  - No definitive recommendation on treatment can be given.
    - **Give NSAID does not work – TRY ANOTHER**



17

### 2019 Aspirin Recommendations

- American College of Cardiology:
  - Aspirin should be used infrequently in the routine primary prevention of ASCVD because of lack of net benefit.
  - low-dose aspirin should not be routinely given as a preventive measure to adults 70 years and older or to any adult who has an increased risk of bleeding
    - Just updated (2021): no longer recommended for patients 60 and older, and for patients aged 40-59 it should be on a case to case basis.



18

## OTC NSAIDs

- **Ibuprofen:**
  - Adult analgesic dose: **200-400mg q4hours**
  - Maximum Dosage: 1200 mg/day OTC for pain (approved for 3200 mg/day in arthritis treatment)
  - OTC: 200 mg tabs (US) 400 mg and 600 mg (Canada)
  - **Most renal toxic of all the NSAIDs**
- **Naproxen sodium:**
  - OTC: 220 mg (Aleve®)
  - **OTC: 1 tablet every 8-12 hours (can use two tablets on first dose)**



19

## Prescription NSAIDs

- **Indomethacin:**
  - Adult Dosage: 25-50 mg TID
- **Mainly used as a short-term anti-inflammatory especially for conditions that do not respond to less toxic NSAIDs.**
  - Indomethacin has a very high level of intolerance compared to other NSAID's.
  - Oral NSAID most widely used in Tx of ocular inflammation.
- **Celebrex:**
  - It is approved for the treatment of osteoarthritis and rheumatoid arthritis.
  - **Dosage: 100 mg BID or 200 mg daily**



20

## Contraindications to NSAIDs

- Avoid in:
  - Pregnancy (especially the late trimesters)
  - **Active Peptic Ulcer Disease**
  - Cross Sensitivity to ASA
  - Previous Hypersensitivity to NSAIDs
  - **Chronic Renal Insufficiency**
- At Risk Patients Include:
  - Dehydration
  - HTN or CHF
    - Use of ACE inhibitors, diuretics and B-blockers
    - **Higher doses of NSAIDs and chronic therapy extending beyond a week will be more likely to increase risk**
  - Advanced Age



21

## NSAID-related ulcers

- COX-2 inhibitors such as celecoxib (Celebrex) are less likely to cause ulcers than aspirin
- **Proton pump inhibitors (e.g. Losec<sup>®</sup>, Prevacid<sup>®</sup> or Prilosec<sup>®</sup>) help to offset the risk of NSAID-related stomach ulcers**
  - patients should be treated with concomitant proton pump inhibitors once daily, which results in ulcer healing rates of approximately 80% at 8 weeks in patients continuing to take NSAIDs



22

## Acetaminophen



- Mechanism of Action is not well understood.
  - Possibly some CNS component
  - Very weak inhibitor of prostaglandin synthesis
- **One of the most commonly used analgesics for mild to moderate pain.**
  - Equal analgesic properties to ASA unless associated with inflammation, where it is less effective.

Take home: Good for pain; Good for fever;  
No effect on inflammation



23

## Dangers of Acetaminophen

- **Acetaminophen overdose is the leading cause of liver failure in the U.S.**
  - It sends 56,000 people to the emergency room annually and causes approximately 400 deaths yearly.
- Acetaminophen is used in so many products, people are often unaware that they are taking it, leading to more overdoses.
  - Combined with agents to get wide range of symptom coverage.
    - Antihistamines such as diphenhydramine – Tylenol PM
    - Diuretics such as Pyrilamine maleate – Midol Complete
    - Cough Suppressants such as Dextromethorphan - Nyquil



24

Consider Combining APAP with NSAID's for Mild to Moderate Pain Relief

- 1:00 pm: Two 325mg acetaminophen
  - 3:00 pm: Two 200mg Ibuprofen
  - 5:00 pm: Two 325mg acetaminophen
  - 7:00 pm: Two 200mg Ibuprofen
- Alternated every 2 hours while awake
- Each medication is q 4 hours.



25

Oral Analgesics: Guidelines

- Never exceed maximum recommended dosages:
  - ASA: 8 grams/day
  - Acetaminophen: 4 grams/day (newer data suggest should be closer to 3-3.2 grams/day)
  - Ibuprofen: 2400 mg/day OTC and up to 3200 mg/day prescription (for RA)
  - Naproxen: 1250/day
  - Naproxen sodium: 1375/day
  - Codeine: 360 mg/day



26

Gabapentin (Neurontin<sup>R</sup>)

- Classified as an anticonvulsant drug
- Additionally, used in the treatment of patients with chronic pain
- Gabapentin has primarily been studied and found effective for the treatment of postherpetic neuralgia and painful diabetic neuropathy; evidence for efficacy in other types of neuropathic pain is limited
  - The transition of gabapentinoids into a first-line pain medication is in part due to an intentional marketing strategy by the pharmaceutical industry without adequate studies.



27

Gabapentin (Neurontin<sup>R</sup>)

- Treatment with gabapentin should be initiated at a low dose with gradual increases until pain relief or dose-limiting adverse effects are achieved.
- Dosage:
  - Day 1 single 300 mg dose
  - Day 2 600 mg dose
  - Day 3 900 mg dose
  - Can be titrated up all the way to 1800 mg/day



28

Gabapentin (Neurontin<sup>R</sup>)

- Gabapentinoids have significant risks despite their reputation as safe drugs.
  - Central nervous system effects such as sedation, dizziness, gait instability, and feeling intoxicated are quite common; as many as one in three patients taking therapeutic doses will experience dizziness or somnolence



29

Opioids Information

- Drug of first choice for the treatment of severe acute pain.
- Block the body's natural protective mechanism for protecting areas in pain – thus never prescribe unless you know the direct cause of the pain.



30

Opioids vs 1000mg Acetaminophen and 400 mg Ibuprofen

- **Ibuprofen Plus Acetaminophen Equals Opioid Plus Acetaminophen for Acute Severe Extremity Pain.** Am Fam Physician. 2018;97(5):348
- **Effect of a Single Dose of Oral Opioid and Nonopioid Analgesics on Acute Extremity Pain in the Emergency Department: A Randomized Clinical Trial.** JAMA. 2017;318(17):1661-1667.
  - no statistically significant or clinically important differences in pain reduction at 2 hours among single-dose treatment with ibuprofen and acetaminophen or with 3 different opioid and acetaminophen combination analgesics



31

**SIGNS OF AN OPIOID OVERDOSE. B.L.U.E.**

**BREATHING** — Breathing during an overdose is shallow, gurgling, erratic, or completely absent.

**LIPS** — Lips and fingertips are blue, due to decreased oxygen throughout the body.

**UNRESPONSIVE** — The victim will not respond to verbal or physical stimulation.

**EYES** — Pupils are pinpoint, as the opioids constrict the pupils to an unusually small size.

32

Opioid Overdose: Management Naloxone (Narcan<sup>®</sup>)

- Opioid antagonist
- Available routes of administration include IV (preferred), IM, SubQ, and intranasal
- **For the initial treatment of an opioid-associated life-threatening emergency, the American Heart Association recommends, after initiation of CPR, the use of intranasal or IM naloxone with a repeat dose as needed.**
- If there is an initial patient response (ie, purposeful movement, regular breathing, moan or other response) but the patient then stops responding, begin CPR and repeat naloxone dose.
- If no initial response, continue CPR and use AED as appropriate



33

Opioid Overdose: Management Naloxone (Narcan<sup>®</sup>)

- **4 mg (contents of 1 nasal spray) as a single dose in one nostril; may repeat every 2 to 3 minutes in alternating nostrils until medical assistance becomes available**



34

Opioid Side Effects

- Side Effects are very hard to predict because opioids can cause CNS depression or stimulation.
- CNS Side Effects
  - **dizziness, lightheadedness, sedation, and drowsiness are the most common**
  - **Mood elevation (euphoria) and disorientation can occur in some patients**
  - Visual symptoms such as blurry vision, miosis, and diplopia can occur.
- Constipation
  - **Opioids inhibit intestinal tract motility**
- **Respiratory Depression**
  - **Most serious side effect of the opioids**



35

Opioids: Codeine

**Note: Tylenol 3 and 4 no longer available as Brand name**

- **Analgesic effect occurs within 20 minutes of ingestion and reaches a maximum at 1 – 2 hours**
  - Ceiling effect occurs.
- Usually administered in combination with acetaminophen .
  - Tylenol 1 (222): codeine 8 mg, 300 mg acetaminophen and 15 mg caffeine (Canada)
  - **Tylenol 3 = Codeine 30 mg and Acetaminophen 300 mg**
    - **Dosage: 1-2 tablets every 4 hours.**
  - Tylenol 4 = Codeine 60 mg and Acetaminophen 300 mg
    - Dosage: 1 tablet every 4 – 6 hours



36

### Codeine

- Also available as generic with 15, 30, or 60 mg of Codeine with 300 mg of Acet. or elixer of 12 mg codeine + 120 mg Acet. per 5 mL.
  - Elixer can be used in children for pain management if >3 years.
- **Serious side effects:**
  - Respiratory depression: caution in patients with asthma, COPD
  - Caution in patients taking sedative medications (Xanax/Valium), muscle relaxants or other pain medications
  - No alcohol consumption



37

### Opioids: Hydrocodone and Oxycodone

- **Hydrocodone:**
  - **Approximately 6X more potent than codeine**
  - Milder Side Effects than Codeine: Less constipation and sedation.
  - Clinically believed to cause more euphoria than codeine, but this is not backed by clinical studies.
- **Oxycodone:**
  - **Approximately 10-12X more potent than codeine**
  - Lower level of side effects in comparison to morphine, but high level of euphoria produced, thus higher level of abuse risk.



38

### Tramadol


- Central acting narcotic
  - Synthetic analogue of codeine.
  - Binds to mu receptors and inhibits norepinephrine and serotonin reuptake.
  - Potential for abuse is very low, but has occurred.
- Available as 50 mg tablets.
- **Dosage: 50 – 100 mg q4 – 6 hours.**
  - Analgesia occurs after 1 hour.
  - Maximum dose: 400 mg/day



39

### Preseptal Cellulitis

- Infection and inflammation located anterior to the orbital septum and limited to the superficial periorbital tissues and eyelids.
- Usually follows sinus infection or internal hordeolum (possibly trauma)
- Eyelid swelling, redness, ptosis, pain and low grade fever.



40

### Differentiating Orbital vs. Preseptal

FINDING	ORBITAL	PRESEPTAL
Visual Acuity	Decreased	Normal
Proptosis	Marked	Absent
Chemosis and Hyperemia	Marked	Rare/Mild
Pupils	RAPD	Normal
Pain and Motility	Restricted and Painful	Normal
IOP		Normal
Temperature	102 - 104	Normal/mild elevation
HA and Assoc. Symptoms	Common	Absent

Treatment: Orals for Preseptal, Often IV for Orbital



41

### Preseptal Cellulitis

- Tx:
  - **Augmentin 500 mg TID or 875 mg BID for 5-7 days**
  - **Keflex 500 mg QID 5-7 days**
  - or if moderate to severe IV Fortaz (ceftriaxime) 1-2 g q8h.
  - if MRSA possible, consider Bactrim/Septa



42

## Antibiotic Resistance

- Microorganism that was originally in the spectrum of activity is no longer susceptible to the drug.
- **Mechanisms of Resistance Include:**
  - Producing an enzyme capable of destroying or inactivating the antibiotic.
  - Altering the target site receptor for the antibiotic so as to reduce or block its binding.
  - Preventing the entry of the antibiotic into the bacterial cell or actively transporting the antibiotic out.



43

## Avoiding Resistance

- Bacterial resistance is a natural result of mutation.
- **Antibiotics cause a faster rate of selection against these resistant bacteria if not prescribed correctly.**
  - Avoid prescribing for non-bacterial infections.
  - Avoid sublethal doses (attack to kill all).
  - Avoid intermittent use.
  - Always complete the full dosage for an appropriate length of time.
  - NEVER TAPER AN ANTIBIOTIC below recommended dosing schedule!



44

## Preventing Resistance

- **The IDSA suggests five to seven days is long enough to treat a bacterial infection without encouraging resistance in adults, though children should still get the longer course**
  - this is different than previous guidelines of treating infections from 10-14 days.

8/6/2024



45

## Antibiotic Associated Diarrhea (AAD)

- The most common side effects of antibiotics are gastro-intestinal, such as nausea and diarrhea
- **AAD arises when the antibiotic disrupts the ecology of the intestinal microbiota, by altering the diversity and numbers of bacteria in the gut.**
- **Diarrhea is most frequently associated with the use of broad-spectrum antibiotics (e.g amoxicillin)**

Agamemnon, V., Kivi, C.A.M., Rijkers, G. et al. A practical guide for probiotics applied to the case of antibiotic-associated diarrhea in The Netherlands. BMC Gastroenterol 18, 103 (2018)



46

## AAD and Probiotics

- **The core benefit of probiotics is exercised by contributing to the maintenance of a balanced microbiota and therefore by creating a favorable gut environment**
- The efficacy of probiotics in preventing AAD depends on the dose.
  - **A daily intake of at least  $5 \times 10^9$  CFU is associated with significant efficacy for AAD** and it has been shown that higher probiotic dose is linked to greater efficacy
  - Example: The probiotic content of yogurt products can range from 90 to 500 billion CFU per serving

Agamemnon, V., Kivi, C.A.M., Rijkers, G. et al. A practical guide for probiotics applied to the case of antibiotic-associated diarrhea in The Netherlands. BMC Gastroenterol 18, 103 (2018)



47

## 5 Facts About Penicillin Allergy

- Approximately 10% of all U.S. patients report having an allergic reaction to a penicillin class antibiotic in their past.
  - **When evaluated, fewer than 1% of the population are truly allergic to penicillins.**
- **Approximately 80% of patients with IgE-mediated penicillin allergy lose their sensitivity after 10 years.**
- Broad-spectrum antibiotics are often used as an alternative to penicillins. The use of broad-spectrum antibiotics in patients labeled "penicillin-allergic" is associated with higher healthcare costs, increased risk for antibiotic resistance, and suboptimal antibiotic therapy.
- **Correctly identifying those who are not truly penicillin-allergic can decrease unnecessary use of broad-spectrum antibiotics.**

<https://www.cdc.gov/antibiotic-use/community/pdfs/penicillin-factsheet.pdf>



48



## Penicillins: Augmentin

- **Augmentin is amoxicillin with potassium clavulanate (clavulanic acid 125 mg).**
- Clavulanate is a B-Lactamase inhibitor which reduces a bacteria's ability to negate the effect of the amoxicillin by inactivating penicillinase (enzyme that inactivates the antibiotic affect).
  - Dicloxacillin can also be used in infections due to penicillinase-producing staph.



49

## Penicillins: Augmentin

- **Augmentin is very effective for skin and skin structure infections such as:**
  - dacryocystitis,
  - internal hordeola,
  - preseptal cellulitis.
- **Treatment of:**
  - otitis media,
  - sinusitis,
  - lower respiratory and urinary infections.
- Given prophylactically to dental surgery patients.



50

## Penicillins: Augmentin

- It has **low:**
  - GI upset,
  - allergic reaction and anaphylaxis.
- **Serious complications include:**
  - anemia,
  - pseudomembranous colitis and
  - Stevens-Johnson syndrome.



51

## Penicillins: Augmentin.

### Adults:

- 250 TID, 500 mg tab BID-TID depending on **what you are treating** (also available in chewable tablets and suspension)
- or 875 mg q 12hr (bid)
- 1000 mg XR: q12 hr and not for use in children <16

### Peds: <3 mos 30mg/kg/day divided q12hrs using suspension

- >3 mos 45-90mg/kg/day divided q12hrs (otitis media 90mg for 10 days)



52

## Cephalosporins

- Closely related structurally and functionally to the penicillins,
  - **have the same mode of action,**
  - affected by the same resistance mechanisms.
  - **tend to be more resistant to B-lactamases.**
- classified as 1st, 2nd, 3rd, 4<sup>th</sup> and now 5th generation based largely on their bacterial susceptibility patterns and resistance to B-lactamases.
- Typically administered IV or IM, **poor oral absorption.**



53

## Side Effects and Contraindications

- **Hypersensitivity Reactions are common.**
  - Risk of cross sensitivity with PCN's is higher for 1<sup>st</sup> generation, but often overestimated for later medications.
  - Used to state the cross sensitivity was ~10%, but now believed to be closer to 3%.



54

## Cephalosporins

- 1st generation: cefadroxil (Duricef), cefazolin (Ancef), **cephalexin (Keflex)**, and cephalothin
- 2nd generations: **cefaclor (Ceclor)**, cefprozil, cefuroxime (Zinacef), cefotetan, cefoxitin
- 3rd generation: **cefdinir (Omnicef)**, cefixime, cefotaxime (Claforan), **ceftazidime (Fortaz)**, ceftibuten, ceftizoxime, ceftriaxone (Rocephin IM/IV).
- 4th generation: cefepime
- 5<sup>th</sup> generation: Cefaroline is a novel **fifth-generation cephalosporin**, which exhibits broad-spectrum activity against Gram-positive bacteria, including MRSA and extensively-resistant strains, such as vancomycin-intermediate *S. aureus* (VISA), heteroresistant VISA (hVISA), and vancomycin-resistant *S. aureus* (VRSA)
- **Keflex, Ceclor, Omnicef** (all orally administered) are effective against most gram positive pathogens and especially good for skin and soft tissue infections.



55

## Cephalosporins

- **Keflex (cephalexin):**
  - treatment of respiratory, GI, skin and skin structure, and bone infections as well as otitis media
  - Adults: 250-1000 mg every 6 hours
    - typical dosing 500 every 6 hours
  - Children: 25-100 mg/kg/day divided 6-8 hours



56

## Cephalosporins

- Cefaclor (Ceclor) (2<sup>nd</sup> generation):
  - Immediate-release: 250 to 500 mg every 8 hours
  - Extended-release: 500 mg every 12 hours

**Note:** An extended-release tablet dose of 500 mg twice daily is clinically equivalent to an immediate-release capsule dose of 250 mg 3 times daily; an extended-release tablet dose of 500 mg twice daily is **NOT** clinically equivalent to 500 mg 3 times daily of other cefaclor formulations.



57

## Cephalosporins

- **Cefdinir (Omnicef no longer available in the US, available as generic)**
  - Used in the treatment of community acquired pneumonia, acute flare ups of chronic bronchitis, acute maxillary sinusitis and tonsillitis.
  - Adult dosing:
    - comes in 300 mg capsules and recommended dosing is 600 mg per day (single dose is equivalent to 300 every 12 hours)



58

## MRSA

- Healthcare-associated methicillin-resistant *Staphylococcus aureus* (HA-MRSA) is associated with severe, invasive disease in hospitalized patients
- Community-associated methicillin-resistant *S. aureus* (CA-MRSA) is most often associated with skin and soft tissue infections in young, healthy individuals with no recent healthcare exposure



59

## Consider Covering for MRSA

1. Hx of non-response to amoxicillin or Augmentin
2. Hx of previous MRSA infections
3. Infection did not start at lid margin like a regular hordeolum but more superior like near the eyebrow area
4. Hx of recent incarceration or hospitalization or in nursing home
5. health care worker
6. pain outside clinical presentation



60

60

Co-Trimoxazole (Bactrim/Septra)

- Combination of trimethoprim and sulfamethoxazole
  - shows greater antimicrobial activity than equivalent quantities of either drug alone.
- Has broader spectrum of action than the sulfa's and is effective in treating:
  - UTIs and respiratory tract infections
  - often considered for treatment of MRSA skin infections



61

61

Co-Trimoxazole (Bactrim/Septra)

- Available:
  - **Bactrim/Septra tablets:**
    - contains 80 mg trimethoprim and 400 mg sulfamethoxazole
    - dosing 2 tablets every 12 hours
  - **Bactrim DS/Septra DS (Double Strength)**
    - contains 160 mg trimethoprim and 800 mg sulfamethoxazole
    - Dosing 1 tablet every 12 hours



62


Case Example

- 67 YOF
- HA and vision loss x 2 days
- OHx: unremarkable
- LEE: 3 days ago!
- MHx: unremarkable


Case courtesy of Dr. Tammy Than



63



64



Case courtesy of Dr. Tammy Than



65

Minocycline?

- Proposed mechanisms
  - ↓ MMPs (MMP-9)
    - Increase in MMP-9 disrupt blood brain barrier and are linked to poor functional recovery
  - Anti-inflammatory
  - Reduction in microglial activation
    - microglial activation is believed to play a central role in neuroinflammation and pathological progression of ischemic tissue
  - Nitric oxide (NO) production
    - NO plays a neuroprotective role in acute ischemic stroke.
  - Inhibition of apoptotic cell death
    - Apoptosis may contribute to a significant proportion of neuron death following acute brain ischemia



66

## Acute Stroke Management

- N=152
- Open-label, evaluator masked study
- Minocycline 200 mg QD x 5 d or placebo
- Evaluated on NIH Stroke Scale
  - 0-1 complete/nearly complete improvement
  - 2-7 – mild
  - 8-14 – moderate
  - >15 – severe
  - Day 30: 1.8 versus 7.1

Total NIH Stroke Scale Score	
1a - Level of Consciousness	1
1b - LOC Questions	1
1c - LOC Commands	1
2 - Best Gaze	0
3 - Visual Fields	0
4 - Facial Palsy	2
5a - Left Motor Arm	0
5b - Right Motor Arm	0
6a - Left Motor Leg	1
6b - Right Motor Leg	0
7 - Language	0
8 - Sensory	1
9 - Best Language	0
10 - Dysarthria	1
11 - Extinction and Intermission	0
Total (NIHSS) Score: 10	

Lamp Y, Boaz M, Giladi R, Lorberboym M, Dabby R, Rasport A, et al. Minocycline treatment in acute stroke. *Neurology*. 2007;69(14):1404-10

TEST	Admission	Day 7	Day 30	Day 90
NHSS - Min	7.5	6.5	1.8	1.6
NHSS - Cont	7.6	8.1	7.3	6.5
mRS - Min	2.8	1.5	1.1	0.9
mRS - Cont	2.0	3.1	2.7	2.1
BI - Min	70.0	85.9	90.6	94.9
BI - Cont	69.9	61.9	68.5	77.6


Minocycline for acute stroke treatment: a systematic review and meta-analysis of randomized clinical trials. *J Neurol*. 2018 Aug;265(8):1871-1879

67

68


## Case Report

- 77 YOM
- Right occipital infarct
- 3 weeks post stroke
  - Minocycline 100 mg BID x 5 days




Mark Tomsik, OD and Marlene Skulskie, OD

Shortly after TX



1 Year Later



69

70

## Tetracyclines

- This group includes:
  - Tetracycline (250mg - 500 mg cap BID-QID) needs to be taken 1 hour before or 2 hours after a meal.
  - Minocycline (100 mg cap BID)
  - Doxycycline (20mg - 100 mg cap or tab BID)
    - In Canada: Apprilon (30 mg doxy + 10 mg slow release doxy)
- Rules of Thumb with Doxy:
  - Do not take before lying down (>2 hours before)
  - Do not take with calcium and avoid antacids
  - Do not take with dairy
  - Do take with food
  - Do educate on sun protection

## Side Effects of Tetracyclines

- Side effects include gastric discomfort, phototoxicity, effects on calcified tissues, vestibular problems, pseudotumor.
- **Pregnancy Category D.**
  - Tetracyclines are attracted to embryonic and growing bone tissue.
    - Depress growth of long bones in pregnant women/children.
    - Cause changes in both deciduous and permanent teeth during the time of tooth development (includes discoloration and increased cavities)
- **Contraindicated in:**
  - Women in the last half of pregnancy
  - Lactating women
  - Children under 8 years of age




71

72

## Meibomian Gland Dysfunction

- Meibomian gland dysfunction:
  - also referred to as meibomitis and patients experience dry eye problems secondary to increased evaporation of the tears.
  - signs include noticeable capping of the glands and frothing of tear film.
- Standard treatment includes:
  - good lid hygiene with warm compresses and lid scrubs in conjunction with
  - **doxycycline 50 mg po BID for 2-3 months**
- **Alternative treatment:**
  - **Azithromycin 500 mg/day for 3 days for three- four weeks**
    - Recent study used single Z-pak treatment



73

## Herpes Zoster

1. Primary infection – Chicken pox (Varicella)
  - Usually in children
  - Highly contagious\*\*\*
  - Very itchy maculopapular rash with vesicles that crust over after ≈ 5 days
  - 96% of people develop by 20 years of age
  - Vaccine now available



74

## Herpes Zoster

### Reactivation – Shingles (Herpes Zoster)


- More often in the elderly and immunosuppressed (AIDS)
  - Systemic work-up if Zoster in someone < 40
- Can get shingles anywhere on the body
- Herpes Zoster Ophthalmicus (HZO)
  - Shingles involving the dermatome supplied by the ophthalmic division of the CNV (trigeminal)
    - 15% of zoster cases




75

## Herpes Zoster

- Associated factors include increasing age, immune deficiency and stress.
- Traditionally thought to only affect patients over the age of 60 and those patients under 60 should be worked up for immune deficiency
- Increasing trend to affect patients of younger age who are not immunocompromised



Shingles and Postherpetic Neuralgia Rates\* by Age, United States



Shingles Rates in Adults 30 and Older, 1998-2016



76

## Herpes Zoster

- Symptoms:
  - Generalized malaise, tiredness, fever
  - Headache, tenderness, paresthesias (tingling), and pain on one side of the scalp
    - Will often precede rash
  - Rash on one side of the forehead
  - Red eye
  - Eye pain & light sensitivity



77

## Herpes Zoster

- Signs:
  - Maculopapular rash -> vesicles -> pustules -> crusting on the forehead
  - Respects the midline\*\*\*
  - Hutchinson sign
    - rash on the tip or side of the nose\*\*\*
  - Classically does not involve the lower lid
  - Numerous other ocular signs



78

## Herpes Zoster

- Other Eye Complications (Acute):
  - Anterior uveitis (most common ocular manifestation)
  - Acute epithelial keratitis (pseudodendrites)
  - Conjunctivitis
  - Stromal (interstitial) interstitial keratitis
  - Endotheliitis (disciform keratitis)
  - Neurotrophic keratitis



79

## Herpes Zoster

- Associated factors include increasing age, immune deficiency and stress.
- Only people who had natural infection with wild-type VZV or had varicella vaccination can develop herpes zoster.
- Children who get the varicella vaccine appear to have a lower risk of herpes zoster compared with people who were infected with wild-type VZV.



80

## Herpes Zoster

- A person's risk for herpes zoster increases sharply after 50 years of age.
- Almost 1 out of 3 people in the United States will develop herpes zoster during their lifetime.
- A person's risk of developing post-herpetic neuralgia also increases sharply with age.



81

## Herpes Zoster

- Management includes:
  - oral antivirals:
    - 800mg acyclovir 5x/day
    - valacyclovir (Valtrex) 1g TID,
    - famciclovir (Famvir) 500 mg TID
  - effectiveness of therapy is best started within 72 hours
  - oral steroids (clinical trials show variable results but often prescribed with antiviral to reduce pain)
  - management of pain (capsaicin, tricyclic antidepressants, gabapentin).
  - If ocular complications, consider topical steroids (Pred Forte QID).



82

## NEW!! Shingrix HZ Vaccine

- Approved in US/Canada as of October 2017
- non-live antigen, to trigger a targeted immune response, with a specifically designed adjuvant to enhance this response and help address the natural age-related decline of the immune system
- Shingrix is 97% effective against shingles for people between the ages of 50 and 69 and 91% effective for people 70 or older.
- It is 91% effective against postherpetic neuralgia for people 50 and older.
- These rates are based on evidence presented to the committee from clinical trials with over 38,000 total participants.



83

## NEW!! Shingrix HZ Vaccine

- recommended for healthy adults aged 50 years and older to prevent shingles and related complications
- recommended for adults who previously received the current shingles vaccine ([Zostavax®](#)) to prevent shingles and related complications
- the preferred vaccine for preventing shingles and related complications



84